


Response to “Comment on ‘Application of an *in Vitro* Assay to Identify Chemicals That Increase Estradiol and Progesterone Synthesis and Are Potential Breast Cancer Risk Factors’”

Ruthann A. Rudel,¹  Bethsaida Cardona,¹ Alexandre Borrel,¹ and Jennifer E. Kay¹

¹Silent Spring Institute, Newton, Massachusetts, USA

<https://doi.org/10.1289/EHP11400>

Refers to <https://doi.org/10.1289/EHP11083>

We appreciate the points raised by Burgoon and Borgert regarding our recent analysis of the U.S. Environmental Protection Agency’s (EPA) data from the high-throughput (HT) H295R screening of 656 chemicals in concentration-response.¹ We considered these same points as we conducted our initial work and further analyzed this issue (complete details are available at https://github.com/SilentSpringInstitute/E2upP4up_followup_Burgoon). We disagree with their assertion that the 296 chemicals we found to increase estradiol (E2) and progesterone (P4) in the HT-H295R assay are mostly false positives.

In order to reach their conclusion about false positives, Burgoon and Borgert made two decisions that are inconsistent with standard approaches used by the U.S. EPA Endocrine Disruptor Screening Program² and the Organisation for Economic Co-operation and Development.³ First, they did not normalize hormone concentration changes from treatment to concurrent plate-specific controls. Second, they required a hormone concentration to increase above a threshold they selected for the response to be positive. Neither the pooling of controls nor the application of a hormone concentration threshold is disclosed in their letter but are described in their GitHub code (https://github.com/DataSciBurgoon/toxcast_steroidogenesis/).

Burgoon and Borgert miscalculated the fraction of positives among all the chemicals that the U.S. EPA screened for two reasons. First, they excluded 84 chemicals tested in 2017, which we included, inflating the fraction positive. In addition, over 2,012 chemicals were initially tested at a single dose in HT-H295R, and 656 were selected for subsequent testing in concentration-response format because they affected multiple hormones. Thus, the 296 positive chemicals we identified as active are derived from screening 2,012 chemicals.

In response to comments from Burgoon and Borgert, we assessed reproducibility of the data by analyzing concordance among the 107 chemicals tested in replicate. We found many

replicates (12 of 23 E2-up chemicals and 18 of 21 P4-up chemicals) showed concordant positive results. Lower concordance among E2-up chemicals is consistent with the overall lower dynamic range of E2 concentration changes in the assay.¹ Overall reproducibility among replicates—especially for P4—adds to the evidence that the data are reliable and that the standard practice of normalization to concurrent controls provides robust results.

Contrary to Burgoon and Borgert’s assertions, we used raw data from the U.S. EPA’s ToxCast database, although—as we made clear in our paper—we used the analysis of variance–based method described by Haggard et al.⁴ to determine the chemical–hormone hit–call instead of using the ToxCast automatic data processing pipeline.⁵ In addition, we mapped the chemicals we identified as increasing E2 or P4 onto data from the National Toxicology Program’s Integrated Chemical Environment database, as Burgoon and Borgert suggested. After filtering for the HT-H295R assay, we found quality control flags for 22 chemicals (13 that increased E2, 4 that increased P4, and 5 that increased both). Most of the flags note low chemical concentration in the well, which can result in false negatives, but positive results are still qualitatively useful.

It is unclear why Burgoon and Borgert highlight “day-effects” only for P4 data and suggest that this is grounds for dismissing the P4 results entirely. Dimethyl sulfoxide controls are included on every plate to account for variability between measurements from different plates. In addition, good concordance between replicates for P4 indicates that this is not a major concern, and P4 results are even more reliable than those for E2.

Acknowledgments

Although our original analysis relied on hit–calls from the U.S. Environmental Protection Agency’s (EPA) analyses of these data,^{4,6} A. Borrel and J. Kay assisted us in conducting additional data analysis to respond to Burgoon and Borgert’s comments, and so we have included them as authors in this response. We also appreciate the support from R.W. Setzer (emeritus U.S. EPA) and K. Paul-Friedman (U.S. EPA) in preparing this response.

References

1. Cardona B, Rudel RA. 2021. Application of an *in vitro* assay to identify chemicals that increase estradiol and progesterone synthesis and are potential breast cancer risk factors. *Environ Health Perspect* 129(7):077003, PMID: [64287026](https://pubmed.ncbi.nlm.nih.gov/34287026/), <https://doi.org/10.1289/EHP8608>.
2. U.S. EPA (U.S. Environmental Protection Agency). 2011. *Steroidogenesis (Human Cell Line – H295R). OCSP Guideline 890.1550. Standard Evaluation Procedure (SEP)*. https://www.epa.gov/sites/default/files/2015-07/documents/final_890.1550_steroidogenesis_assay_sep_8.1.11.pdf [accessed 7 April 2022].
3. Organisation for Economic Co-operation and Development. 2011. Test No. 456: H295R Steroidogenesis Assay. OECD Guidelines for the Testing of Chemicals, Section 4. <https://doi.org/10.1787/9789264122642-en>.
4. Haggard DE, Karmaus AL, Martin MT, Judson RS, Setzer RW, Paul Friedman K. 2018. High-throughput H295R steroidogenesis assay: utility as an alternative and a statistical approach to characterize effects on steroidogenesis.

Address correspondence to Ruthann A. Rudel, Silent Spring Institute, 320 Nevada St., Newton, MA 02460 USA. Email: Rudel@SilentSpring.org

This work was funded by the U.S. National Institutes of Health/National Institute of Environmental Health Sciences Breast Cancer and Environment Research Program (U01ES026130), the California Breast Cancer Research Program (awards 21UB-8012 and 21UB-8100), the Cedar Tree Foundation, and charitable gifts to the Silent Spring Institute. All authors are or have been employed at the Silent Spring Institute, a scientific research organization dedicated to studying environmental factors in women’s health. The Silent Spring Institute is a 501(c)3 public charity funded by federal grants and contracts, foundation grants, and private donations, including from breast cancer organizations. Study funders had no role in the study design; in the collection, analysis, or interpretation of data; in the writing of the report; or in the decision to submit the article for publication.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehpsubmissions@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

- Toxicol Sci 162(2):509–534, PMID: [29216406](https://pubmed.ncbi.nlm.nih.gov/29216406/), <https://doi.org/10.1093/toxsci/kfx274>.
5. Filer DL, Kothiya P, Setzer RW, Judson RS, Martin MT. 2017. tcpl: the ToxCast pipeline for high-throughput screening data. *Bioinformatics* 33(4):618–620, PMID: [27797781](https://pubmed.ncbi.nlm.nih.gov/27797781/), <https://doi.org/10.1093/bioinformatics/btw680>.
6. Haggard DE, Setzer RW, Judson RS, Paul Friedman K. 2019. Development of a prioritization method for chemical-mediated effects on steroidogenesis using an integrated statistical analysis of high-throughput H295R data. *Regul Toxicol Pharmacol* 109:104510, PMID: [31676319](https://pubmed.ncbi.nlm.nih.gov/31676319/), <https://doi.org/10.1016/j.yrtph.2019.104510>.